



## Hennepin County Adolescent Sexual Health Report

### Sexual Activity

According to the 2001 Minnesota Student Survey, in Hennepin County, 14% of 9th grade females and 22% of 9th grade males were sexually active; 42% of 12th grade females and 43% of 12th grade males were sexually active.<sup>i & ii</sup>

Additionally, 47% of sexually active 9<sup>th</sup> grade females, 41% of 9<sup>th</sup> grade males, 68% of 12<sup>th</sup> grade females, and 64% of 12<sup>th</sup> grade males reported always using contraception. 34% of sexually active 9<sup>th</sup> grade females, 38% of 9<sup>th</sup> grade males, 15% of 12<sup>th</sup> grade females and 18% of 12<sup>th</sup> grade males reported never or rarely using a method of contraception.<sup>iii</sup>

<b>Percentage of sexually active teens:</b>		
	<b><u>Female</u></b>	<b><u>Male</u></b>
<b>9th grade</b>	14%	22%
<b>12th grade</b>	42%	43%

### Pregnancy and Births

In Hennepin County in 2000, 699 teens aged 15-17 years old became pregnant, and 1373 teens aged 18-19 years old became pregnant.<sup>iv</sup> In total, 2072 teens aged 15-19 years old became pregnant. The combined 1998-2000<sup>v</sup> pregnancy rate for 15-17 year olds was 34.8, for 18-19 year olds was 91.2, and for 15-19 year olds was 58.2.<sup>vi</sup>

In 2000, there were 32 births to females under 15 years old in Hennepin, 458 births to 15-17 year olds, and 842 births to 18-19 year olds. In total, there were 1300 births to 15-19 year olds. The combined 1998-2000 birth rate for 15-17 year olds was 22.2, for 18-19 year olds was 55.3, and for 15-19 year olds was 35.9.<sup>vii</sup>

<b>Number of Pregnancies 2000:</b>	
<b>15-17 years old</b>	699
<b>18-19 years old</b>	1373
<b>15-19 years old</b>	2072

<b>Pregnancy Rates 1998-2000</b> (per 1000 females in each age group)	
<b>15-17 years old</b>	34.8
<b>18-19 years old</b>	91.2
<b>15-19 years old</b>	58.2

<b>Number of Births 2000:</b>	
<b>Under 15 years old</b>	32
<b>15-17 years old</b>	458
<b>18-19 years old</b>	842
<b>15-19 years old</b>	1300

<b>Birth Rates 1998-2000:</b> (per 1000 females in each age group)	
15-17 years old	22.2
18-19 years old	55.3
15-19 years old	35.9

### STDs and HIV

In Hennepin County, 467 teens aged 15-19 years old were diagnosed with gonorrhea in 2000.<sup>viii</sup> 1073 teens, aged 15-19 years old were diagnosed with chlamydia in 2000. According to the MN Department of Health, in the year 2000 nine HIV/AIDS cases were diagnosed in Minnesota teenagers 15 to 19 years of age. 4 of the nine were diagnosed when they were ages 15-17, 5 of the 9 were diagnosed when they were 18-19 years of age. 6 were female. 3 were male. 4 cases were White/Non-Hispanic, 2 Black/Non-Hispanic, 2 Hispanic, and 1 American Indian/Alaskan Native. 8 of the 9 residents were residents of Hennepin County and 1 was from Anoka County. There are not enough cases to analyze in more detail.

According to the 2001 Minnesota Student Survey, 60% of sexually active 9<sup>th</sup> grade females, 50% of 9<sup>th</sup> grade males, 71% of 12<sup>th</sup> grade females and 61% of 12<sup>th</sup> grade males reported they had spoken with every sexual partner about protecting themselves from STDs. Also, 73% of sexually active 9<sup>th</sup> grade females, 72% of 9<sup>th</sup> grade males, 56% of 12<sup>th</sup> grade females and 65% of 12<sup>th</sup> grade males reported using a condom at last intercourse.

### **Prenatal Care/Low Birth Weight**

Adequate use of prenatal care services is good insurance for a healthy pregnancy, birth and baby. Teens in the United States are less likely to get adequate prenatal care than adult women.<sup>ix</sup> Of those whose prenatal care use was reported to the Minnesota Department of Health in 2000 in Hennepin County, 9% of pregnant women ages 15 to 19 received no care or did not receive care until the third trimester (versus 4% of women ages 20-29 and 2% of women ages 30-39). Additionally, 9% of births to teens ages 15 to 19 in 2000 resulted in infants who were reported as low birth weight (versus 7% of births to women ages 20-29 years old and 7% of women ages 30-39).<sup>x</sup> It should be noted that the association between maternal biological age and low birth weight can sometimes be very strong, but the association does not appear to be causal. Low birth weight is strongly associated with poverty; women who are childbearing as teens are more likely to be poor than women who wait until their 20's or 30's to have children.<sup>xi</sup>

### **Public Assistance**

While few teen parents in Minnesota and in the United States receive public assistance, families that began with a teen giving birth are more likely to be on public assistance than those who first gave birth between 20 and 24 years of age.<sup>xii</sup> Additionally, women who start childbearing in their teen years and who start on public assistance at any time are likely to remain on it longer.<sup>xiii</sup> As of December, 2001, 6678 MFIP (Minnesota Family Investment Program) families, or 55% of all families receiving MFIP in Hennepin County, began with a teen giving birth.<sup>xiv</sup>

**The Minnesota Organization on Adolescent Pregnancy, Prevention and Parenting is a statewide membership organization which provides a newsletter, an annual conference, regional trainings, public policy information, and the InfoExchange, a clearinghouse of information on issues relating to teenage pregnancy prevention and teenage parenting. MOAPPP's mission is to strengthen policies and programming related to adolescent pregnancy, prevention and parenting in Minnesota.**

### **Notes**

<sup>i</sup> Minnesota Student Survey 2001, Minnesota Department of Children, Families and Learning, (651)582-8495.

<sup>ii</sup> Sexually active is defined as having had sexual intercourse one or more times. It does not necessarily mean that the teen is currently having sex.

<sup>iii</sup> Totals may exceed 100% due to rounding.

<sup>iv</sup> Minnesota Department of Health, Minnesota Center for Health Statistics, 2000 birth and pregnancy statistics, (651) 297-1232.

<sup>v</sup> Multiple year averaging of rates is important because it increases the number of "events" (pregnancies and births) being counted. Therefore the more variable one-year rates become less noticeable and the three-year average provides a better reflection of the "true" rate of pregnancies than will three consecutive annual rates.

<sup>vi</sup> Pregnancy rate refers to the number of live births plus the number of fetal deaths plus the number of induced abortions per 1,000 females in the population of the specified age.

<sup>vii</sup> Number of live births per 1,000 population.

<sup>viii</sup> Minnesota Department of Health, (612) 676-5461.

<sup>ix</sup> The Alan Guttmacher Institute. (1994). *Sex and America's Teenagers*. New York.

<sup>x</sup> Low birth weight is defined as less than 2500 grams.

<sup>xi</sup> Chomitz, V.R., Cheung, L.W., Lieberman, E. (1995). "The Role of Lifestyle in Preventing Low Birth Weight." *The Future of Children*, vol.5 (1): 121-138.

<sup>xii</sup> The Alan Guttmacher Institute. (1994). *Sex and America's Teenagers*. New York.

<sup>xiii</sup> *Kids Having Kids*, The Robin Hood Foundation Special Report on the Cost of Adolescent Childbearing, New York, 1996.

<sup>xiv</sup> Minnesota Department of Human Services, Division of Reports and Forecasts; 2001 data, prepared by Paul Farseth, (651) 296 - 8560.